

2009
2009/10/17

**A study of carotenoids of mango fruit and mango based
products and their *in-vitro* bioavailability**

by

Matara Arahchige Jagath wansapala

Libary - USJP



200487

**Thesis submitted to the University of Sri Jayewardenepura for
the awarded of the Degree of Doctor of Philosophy in Food
Science on 2009**

200487

**Study of carotenoids of mango (*Mangifera indica*) fruit and mango based products
and their *in vitro* bioavailability**

By Matara Arahchighe Jagath Wansapala

ABSTRACT

Vitamin A is an important micronutrient of animal origin responsible for structure and functioning of the skin, mucous membrane and maintaining of a good vision. Low-income, poor educational background and cultural and religious practices discourage the intake of recommended amount of vitamin A through animal sources. Therefore the attention should be given to the naturally available and affordable provitamin A sources such as green leafy vegetables, certain root crops and most of the fruits.

The major objectives of this study were to qualitatively analyze the carotenoid profile in three popular mango varieties namely “Karuthacolamban”, “Beti Amba” and “Gira Amba” grown in the three different agro climatic regions of the Sri Lanka, quantification and *in vitro* bioavailability of pro vitamin A content in all three varieties with a view to updating the database. The impact of different process conditions on the stability of provitamin A carotenoids in mango based products (ie. jam, cordial and osmotically dehydrated mango) were also studied. Sensory characteristics of the selected mango varieties were comparatively studied and the sensory web was developed for each variety.

A reversed phase HPLC method has been developed for the study for the separation and extraction of the carotenoids in crude and saponified extracts from the “Karuthacolamban”, “Beti Amba”, “Gira Amba” and mango based products. All fruits with the same maturity

indices were selected from local market and the cultivars were identified by Horticultural Crop Research and Development Institute Gannoruwa, Kandy, Sri Lanka.

Carotenoids from mango were extracted according to the method described by Rodriguez Amaya (1999), which involved extraction of carotenoids, partition to petroleum ether, separation of carotenoids by Open Column Chromatography (OCC), identification of carotenoids using Ultra violet Visible absorption spectra (maximum absorption and spectral fine structure), order of elution in OCC and chemical tests. Purity of the identified carotenoids were further confirmed by using the High Performance Liquid Chromatography with photo diode array detection (C_{18} column Spherisorb ODS2, $5\mu\text{m}$, $4.6\text{mm} \times 150\text{mm}$; gradient elution of mobile phase of Methanol, Acetonitrile and 0.05% Tri ethyl amine in Ethyl acetate). The HPLC analysis of crude extract of variety Karuthacolamban highlighted the existence of four main peaks which had UV- Visible spectra similar to those reported for violaxanthin, neoxanthin, β -carotene and α -cryptoxanthin. The HPLC analysis of crude extract of "Beti Amba" and "Gira Amba" varieties highlighted the existence of three main peaks which had UV- Visible spectra similar to those reported for violaxanthin, neoxanthin, and β -carotene. A saponification step of crude extracts indicated the existence of esterified violaxanthin, neoxanthin, and α -cryptoxanthin. The Variety "Karuthacolomban" contains $2.7 \pm 0.3 \mu\text{g/g}$ (Fresh weight), the variety "Beti Amba" contains $2.6 \pm 0.3 \mu\text{g/g}$ (Fresh weight) and the variety "Gira Amba" ($0.18 \pm 0.6 \mu\text{g/g}$ (Fresh weight) of β - carotene as the principal pro-vitamin A carotenoid. The amount of in vitro bioavailability β - carotene was highest in the "Beti Amba" variety (29.6%) than "Karuthacolomban" (24%) and "Gira Amba" (20.30%).

Thus, this study has indicated that varietal difference does not exist in the content but bioavailability of β -carotene in mango.

The percentages of destruction of β - carotene in jam, cordial and osmotically dehydrated mango during the process were recorded as 89%, 78% and 66% respectively. The in vitro bioaccessibility of β - carotene in all three products were recorded as 67 %, 25 % and 64% respectively. Higher amounts of provitamin A carotenoids (1.43 ± 0.3 $\mu\text{g/ g}$ Fresh weight) were obtained under Room temperature (28 °C) and the lowest amount of provitamin A carotenoids (0.20 ± 0.3 $\mu\text{g/ g}$ Fresh weight) were recorded at -2 °C. The provitamin A carotenoids content of artificially ripened mango was (0.30 ± 0.3 $\mu\text{g/ g}$ Fresh weight) lower than that of the natural. The highest sensory values were recorded “Gira amba” variety.

Table of content

| | | Page |
|-----|--|-----------|
| I | LIST OF TABLES | vi -vii |
| II | LIST OF FIGURES | viii - xi |
| III | ACKNOWLEDGEMENTS | xii |
| IV | ABSTRACT | xiii -xiv |
| | CHAPTER I : INTRODUCTION | 1 |
| | 1.1 Background | 1 |
| | 1.2 Importance of Mango (<i>Mangifera indica</i>) as a source of carotenoids | 3 |
| | 1.3. Objectives of the study | 5 |
| | CHAPTER II : REVIEW OF LITERATURE | 8 |
| | 2.1 Carotenoids | 8 |
| | 2.1.1 Chemistry of carotenoids | 8 |
| | 2.1.2 Stability of carotenoids | 12 |
| | 2.1.3.Mechanism of carotenoid degradation | 13 |
| | 2.1.4.Biosynthetic pathway of carotenoids | 15 |
| | 2.1.5.Factors affecting the carotenoid composition of fruits and vegetables | 18 |
| | 2.2 Impact of carotenoids on human health | 21 |
| | 2.2.1 Vitamin A and its importance on human health | 22 |
| | 2.2.2. Assessment of vitamin A Status | 23 |
| | 2.2.3 Hypervitaminosis A | 26 |
| | 2.2.4 Bioavailability of pro vitamin A carotenoids | 27 |
| | 2.2.4.1 Factors that affect carotenoid bioavailability | 28 |
| | 2.2.5. Pro vitamin A activity of carotenoid | 30 |
| | 2.2.6 Antioxidant functions of carotenoids | 32 |

| | |
|--|-----------|
| 2.2.7. Carotenoids as food colours | 34 |
| 2.3 Mango (<i>Mangifera indica</i> L.) | 37 |
| 2.3.1 Ripening process of mango | 41 |
| 2.3.2 Mango(<i>Mangifera indica</i> L.) carotenoids | 42 |
| 2.3.2.1 Stability of mango carotenoids under different conditions | 44 |
| 2.4 Important carotenoids in vegetables and other common fruits. | 44 |
| 2.4.1 Important carotenoids in green Leafy vegetables. | 44 |
| 2.4.2 Important carotenoids in root crops. | 45 |
| 2.4.3 Important carotenoids in common Sri Lankan fruits other than Mango. | 45 |
| 2.5. Guide line for carotenoid analysis | 46 |
| 2.6. Chemical test for carotenoids | 48 |
| 2.7 Intestinal absorption of dietary carotenoids | 48 |
| 2.7.1 The theory for <i>in vitro</i> bioaccessibility carotenoids. | 49 |
| CHAPTER III :MATERIALS AND METHODS | 52 |
| 3.1 Extraction of carotenoids from three mango varieties (Karuthacolamban, Beti amba and Gira amba) | 52 |
| 3.1.1 Selection of samples | 53 |
| 3.1.2. preparation of samples | 54 |
| 3.1.3. Isolation of carotenoids | 54 |
| 3.2 Saponification of carotenoids | 55 |
| 3.3. Separation and identification of carotenoids | 56 |
| 3.3.1. Separation of carotenoids by Open Column Chromatography (OCC) | 56 |
| 3.3.2. Separation of carotenoids by Thin Layer Chromatography (TLC) | 59 |
| 3.3.3. Separation of carotenoids by High Performance Liquid Chromatography (HPLC) | 60 |
| 3.4. Chemical tests for the positions of substituents of carotenoids | 62 |
| 3.4.1. Acetylation of Primary and Secondary Hydroxyl Group | 63 |

| | |
|--|-----------|
| 3.4.2. Methylation of allylic Hydroxyl Group | 63 |
| 3.4.3 Epoxide- Furanoid rearrangement | 64 |
| 3.4.4 .Reduction of a conjugated carbonyl group | 64 |
| 3.4.5.Iodine catalized cis – trans isomerization | 64 |
| 3.5 Production process of mango based products (i.e. jam , cordial and osmotically dehydrated mango) | 64 |
| 3.5.1. Jam manufacturing process | 64 |
| 3.5.2. Cordial manufacturing process | 67 |
| 3.5.3. Osmotically dehydrated mango manufacturing process | 69 |
| 3.6. <i>In vitro</i> bioaccessibility of pro vitamin A carotinoids | 71 |
| 3.6.1. <i>In vitro</i> bioaccessibility of pro vitamin A carotinoids in selected mango varieties | 73 |
| 3.6.2. <i>In vitro</i> bioaccessibility of pro vitamin A carotinoids in mango based Jam | 74 |
| 3.6.3. <i>In vitro</i> bioaccessibility of pro vitamin A carotinoids in mango based Cordial | 74 |
| 3.6.4. <i>In vitro</i> bioaccessibility of pro vitamin A carotinoids in Osmotically dehydrated mango. | 74 |
| 3.7. Effect of storage conditions on the provitamin A carotenoid profile of Beti amba variety | 75 |
| 3.7.1 Effect of storage temperature on the provitamin A carotenoid profile of Beti amba variety | 75 |
| 3.7.2 Carotenoid profile of Beti amba variety under artificial ripening environment | 76 |
| 3.8. Distribution of carotenoids within the mango flesh | 76 |
| 3.9. Development of sensory web for selected mango varieties | 76 |
| CHAPTER IV :RESULTS | 77 |
| 4.1 Extraction and Isolation of carotenoids (i.e. Karuthacolamban, Beti amba and Gira amba) | 77 |

| | |
|---|------------|
| 4.1.1. Selection of samples | 77 |
| 4.2. Separation and identification of carotenoids (i.e. Karuthacolamban, Beti amba and Gira amba) | 78 |
| 4.2.1. carotenoids profile of variety karuthacolamban | 79 |
| 4.2.2 Carotenoid profile of variety Beti Amba | 83 |
| 4.2.3. Carotenoid profile of variety Gira Amba | 87 |
| 4.3. Carotenoid profile of mango based products | 91 |
| 4.3.1. Carotenoid profile of Jam | 91 |
| 4.3.2. Carotenoid profile of Cordial | 92 |
| 4.3.3. Carotenoid profile of Osmotically dehydrated mango | 93 |
| 4.4 <i>in vitro</i> bioaccessibility of pro vitamin A carotenoids in the mango varieties(i.e. Karuthacolamban , Beti amba and Gira amba) | 94 |
| 4.4.1 Mango variety – Karuthacolamban | 94 |
| 4.4.2 Mango variety – Beti amba | 95 |
| 4.4.3. Mango variety- Gira amba | 95 |
| 4.5 Pro vitamin A carotenoids and their <i>in vitro</i> bioaccessibility of mango based products (i.e. Jam , Cordial and Osmotically dehydrated mango). | 97 |
| 4.5.1. Mango Jam | 98 |
| 4.5.2. Mango Cordial | 98 |
| 4.5.3. Osmotically dehydrated mango | 99 |
| 4.6. Effect of storage conditions on pro vitamin A carotenoids of the selected mango variety. | 100 |
| 4.6.1. Effect of artificial ripening on pro vitamin A carotenoids | 102 |
| 4.7. Results of the sensory evaluation | 103 |
| CHAPTER V :DISCUSSION | 110 |
| CHAPTER VI :CONCLUSION | 131 |
| REFERENCES | 134 |

| | |
|---|------------|
| APPENDIX I- List of publications and communications from the study | 148 |
| APPENDIX II – Spectrums relevant to the major carotenoids | 149 |