

**Assessment of the Genetic Diversity of Coconut with Special Reference
to Phenotypes in the Galle District in Sri Lanka**

By

Gamachchige Kingsly Ekanayake



M. Phil

2010

**Assessment of the Genetic Diversity of Coconut with Special
Reference to Phenotypes in the Galle District in Sri Lanka**

By

Gamachchige Kingsly Ekanayake

**Thesis submitted to the University of Sri Jayawardenepura for
the award of the Degree of Master of Philosophy in Botany on
31st December 2010**

“The work described in this thesis was carried out by me under the supervision of Dr. S.A.C.N. Perera, Dr. P.N. Dassanayaka and Mr. J.M.D.T. Everard and a report on this has not been submitted in whole or in part to any other university or any other institution for another Degree/Diploma.”




G.K. Ekanayake

"We certify that the above statement made by the candidate is true and that this thesis is suitable for submission to the University for the purpose of evaluation"




15.01.2013

Dr. (Mrs) P. N. Dassanayaka
(Internal supervisor)



16.01.2013

Dr. (Ms.) S. A. C. N. Perera
(External supervisor)



16/01/2013

Dr. J. M. D. T. Everard
(External supervisor)

Table of Contents

	Page
Table of contents	i
List of tables	vii
List of figures	ix
List of plates	x
Acknowledgements	xi
Dedication	xii
Abstract	xiii
CHAPTER 1: INTRODUCTION	01
1.1 Coconut	01
1.2 Coconut Genetic Resources in Sri Lanka	02
1.3 Improvement of production potential of coconut	04
1.4 Importance of coconut breeding in Sri Lanka	05
1.5 Conservation and characterization of coconut germplasm in Sri Lanka	07
1.6 Objectives	09
CHAPTER 2: REVIEW OF LITERATURE	10
2.1 Coconut Palm	10
2.1.1 Botanical description	10
I) The leaf	11
II) The root	12
III) The Stem	13
IV) The Inflorescence	13

V) The Female Flower	14
VI) The number of female flowers	15
VII) The Male Flower	16
VIII) The Pollen	16
IX) The Fruit	16
2.2 Evolution, distribution and dispersion of coconut	18
2.2.1 Origin	18
2.2.2 Evolution	18
2.2.3 Distribution	19
2.2.4 Dispersion	19
2.3 Taxonomy and classification of coconut	20
2.3.1 Classification of coconut in Sri Lanka	20
2.4 Coconut Breeding	22
2.4.1 Coconut breeding in Sri Lanka	22
2.4.2 Objectives of coconut breeding in Sri Lanka	23
2.4.3 Achievements in coconut breeding in Sri Lanka	24
2.5 Collecting and Conservation of coconut germplasm	26

2.5.1 Coconut germplasm and coconut breeding	26
2.5.2 Importance and the necessity for conservation of coconut germplasm	26
2.5.3 Collecting and conservating of coconut germplasm in Sri Lanka	27
2.5.4 Coconut genetic resources in Sri Lanka	28
2.6 Characterization of coconut genetic resources	28
2.6.1 Morphological characterization and evaluation	29
2.7 Genetic polymorphisms for crop improvement	31
2.7.1 Genetic Markers	31
2.7.2 Morphological Markers	32
2.7.3 Molecular markers	32
2.7.4 Advantages of Molecular Markers over Morphological Markers	33
2.7.5 DNA markers	34
I) Restriction Fragment Length Polymorphism (RFLP)	34
II) Random Amplified Polymorphic DNA (RAPD)	35
III) Amplified Fragment Length Polymorphism (AFLP)	35
IV) Simple Sequence Repeats – Microsatellites (SSR)	36
V) Single Nucleotide Polymorphisms (SNP)	38
VI) Diversity Arrays Technology (DArT)	38

2.8 Molecular Marker based assessments in coconut	39
2.8.1 Use of molecular markers in germplasm characterization of coconut	40
2.9 Germplasm diversity and evaluation for coconut improvement	46
Chapter 03: Materials and Methods	47
3.1 Germplasm exploration survey	47
3.1.1 Selection of sites	47
3.2 Conservation of selected coconut phenotypes	48
3.2.1 Procedure of self pollination	48
I) Pollen processing	48
II) Pollination	49
3.3 Morphological characterization	49
3.3.1 Material	49
3.3.1 Morphological Descriptors	50
3.3.2 Analysis of morphological data	53
3.4 Molecular characterization	53
3.4.1 Plant Material	53
3.4.2 DNA Extraction	54
3.4.3 Quantification of DNA	56
3.4.4 SSR assay	56
3.4.4.1 Primers	56
3.4.4.2 Polymerase Chain Reaction (PCR)	57

3.4.5 Polyacrylamide gel electrophoresis for detection of SSR Polymorphisms	57
CHAPTER 4: RESULTS AND DISCUSSION	61
4.1 Germplasm exploration programme	61
4.2 Documentation and the conservation of the new coconut phenotypes	62
4.2.1 Visually observed morphological descriptions of new phenotypes	62
4.2.2 <i>Ex-situ</i> Conservation of new coconut phenotypes	68
4.3 Morphological characterization of the coconut phenotypes	68
4.3.1 Qualitative morphological characters	68
I) Varietal Classification of new phenotypes of coconut	69
II) Classification of the new coconut phenotypes into forms within varieties	70
4.3.2 Quantitative morphological characters	72
4.3.2.1 Descriptive statistics and Analysis of Variance	72
4.3.2 Multivariate data analysis	84
I) Leaf Morphology	84
II) Inflorescence morphology	87
III) Fruit morphology	91
IV) Principal Component Analysis for quantitative and qualitative traits	95
4.4 Molecular characterization of the coconut phenotypes	103
4.4.1 Allelic variations and polymorphisms	103
4.4.2 DNA fingerprints for coconut forms and new phenotypes	109

4.4.3 Genetic distances and relatedness of coconut forms and phenotypes	112
4.5: General Discussion	117
4.5.1 Enrichment of the local coconut germplasm pool	117
4.5.2 Morphological variation of the new coconut germplasm	119
4.5.3 Detection of molecular variation at microsatellite marker loci	121
CHAPTER 5: CONCLUSIONS	124
REFERENCES	125
APPENDIX	137

Lists of Tables

	Page
Table 2.1: Coconut varieties and forms in Sri Lanka	21
Table 3.1: Number of palms selfed in each phenotype	48
Table 3.2 Sample size of each phenotype used for morphological characterization	50
Table 3.3: Descriptors used in morphological characterization	51
Table 3.4: No. of replicate palms of the coconut phenotypes used for molecular characterization	54
Table 3.5: Primer sequences and annealing temperatures of the microsatellite markers used for genotyping	60
Table 4.1: Comparison of morphological descriptors of coconut varieties in Sri Lanka with the new phenotypes	69
Table 4.2: Specific morphological features and the suggested classification of the new coconut types	71
Table 4.3: Comparison of the nut size and kernel weights of the dwarfs with relevant <i>Murusi</i> form	72
Table 4.4 Descriptive statistics of stem parameters of the scored phenotypes	74
Table 4.5 Descriptive statistics of leaf characters of the scored phenotypes	76
Table 4.6 Descriptive statistics of inflorescence characters of the scored phenotypes	78

Table 4.7 Descriptive statistics of fruit characters of the scored phenotypes	81
Table 4.8: Eigen-analysis of the correlation matrix for leaf characters	84
Table 4.9 Distance matrix of the coconut phenotypes based on leaf characters	87
Table 4.10: Eigen-analysis of the correlation matrix for inflorescence characters	88
Table 4.11 Distance matrix for inflorescence characters	91
Table 4.12: Eigen-analysis of the correlation matrix for inflorescence characters	92
Table 4.13: Distance matrix for fruit characters	94
Table 4.14: Eigen-analysis of the correlation matrix for inflorescence characters	96
Table 4.15 Distance matrix based on all the characters scored	99
Table 4.16 Pearson correlation matrix of the measured characters	101
Table 4.17 Summary statistics of genotypic data	103
Table 4.18: Allele frequencies observed in the population	107
Table 4.19: Distance matrix based on genotypic data	113

List of Figures

	Page
Figure 4.1: PCA scatter plot for leaf characters of the coconut phenotypes scored	85
Figure 4.2: Dendrogram for leaf characters based on Euclidean distances	86
Figure 4.3: PCO scatter plot for inflorescence characters of the coconut phenotypes scored	89
Figure 4.4 Dendrogram for inflorescence characters based on Euclidean distances	90
Figure 4.5 PCA scatter plot for fruit characters of coconut phenotypes	93
Figure 4.6: Dendrogram for fruit characters based on Euclidean distances	94
Figure 4.7: PCA scatter plot for all the characters measured in coconut phenotypes	97
Figure 4.8: Dendrogram for all characters based on Euclidean distances	98
Figure 4.9 Graphical representation of the genotypes of each individual	111
Figure 4.10: Phenetic tree drawn based on shared allele distances	116

List of Plates

	Page
Plate 4.1: Ran pol	66
Plate 4.2: Juwan coconuts	66
Plate 4.3: Green, yellow and brown Murusi	66
Plate 4.4: Bothal thembili	67
Plate 4.5 Thatin coconut	67
Plate: 4.6 Rath gon thembili	67
Plate 4.7: Dothalu coconut	67
Plate 4.8: Carpled coconuts	67
Plate 4.9: SSR gel image for the marker CNZ-44 of the forty coconut individuals genotyped	105
Plate 4.10: SSR gel image for the marker CNZ-46 of the forty coconut individuals genotyped	105

Acknowledgements

I would like to express my deepest gratitude to my supervisor Dr. (Ms) S.A.C.N. Perera, Principal Research Officer of the Genetics & Plant Breeding Division of the Coconut Research Institute of Sri Lanka, for her invaluable guidance, immeasurable support and cooperation provided to make this project a success. I am also greatly indebted to my supervisors, Dr (Mrs.) Nilanthie Dassanayake, Senior Lecturer of the University of Sri Jayawardenepura and Dr. J.M.D.T. Everard, the Deputy Director (Research) of the Coconut Research Institute of Sri Lanka for their valuable guidance, immense support and important advices kindly provided throughout this project.

I am grateful to Dr. (Mrs.) C. Jayasekera, the Director of the Coconut Research Institute of Sri Lanka for granting permission to carry out this study as my M. Phil research. I am also very much grateful to Dr. Lalith Perera, Head of the Genetics and Plant Breeding Division of CRI and Dr. (Mrs.) Champa Bandaranayake, former Senior Research Officer of the same division for their invaluable inputs provided for this research.

I would like to express my sincere gratitude to Mrs. Sandya Fernando, Mr. H.M.N.B. Herath, Mr. Ayodya Fernando, Miss Auchithya Dissanayake, Mr. J.H. Somaratne, Mr. P.A.D. Milton, Mrs Indira Jayawardene and all the other staff members of the Genetics & Plant Breeding Division for their continuous support in various ways to make this research a success. A special word of thanks goes to Mr. Athula Senarath and Mr. Priyantha Kumara attached to the Genetics and Plant Breeding Division of CRI for their help in this research.

I would like to acknowledge the financial assistance of the Coconut Genetic Resources Network for farmers' variety survey and the Coconut CESS fund for characterization.

I would like to express my gratitude to Mrs. Srimalie Mallawaarachchi and Ms. T.D.S. Padmini for their valuable assistance provided in various ways to make this project a success. Finally, I express my heartfelt gratitude to my parents for their immeasurable love and care bestowed to me throughout my life.

This thesis is dedicated to my loving parents

**Assessment of the Genetic Diversity of Coconut with Special Reference to
Phenotypes in the Galle District in Sri Lanka**

G. K. Ekanayake

ABSTRACT

Genetic base of the coconut germplasm in Sri Lanka has been reported to be rather narrow. This has posed a serious restriction on the effort of the coconut breeders in developing new coconut cultivars which are essential in fulfilling the objectives of coconut breeding. In an effort to overcome this problem Coconut Research Institute of Sri Lanka initiated a programme for the identification of less known, diverse farmers' varieties of coconut. The research described in this thesis was carried out with the objective of identification of the farmers' varieties of coconut, *ex-situ* conservation of the newly identified material and *in-situ* characterization of the new phenotypes with morphological and molecular methods. Out of the three locations; Wilpotha, Hettipola and Unawatuna where the germplasm exploration programme was carried out, the latter area in the Southern province proved to have diverse coconut germplasm which have not been identified in the previous Sri Lankan coconut classifications. Consequently, several new coconut phenotypes with specific vernacular names such as Ran pol, Juwan pol, Bothal thembili, Murusi, Thatin, Dothalu *etc* and several other phenotypes with no such vernacular names were identified and documented. A pollination programme was carried out to multiply such material and the resultant seedlings were conserved in the *ex-situ* field gene banks of coconut belonging to the Coconut Research Institute of Sri Lanka.

IPGRI descriptors for coconut were used to *in-situ* characterize the morphological diversity, of the new coconut phenotypes along with comparative already classified coconut forms, Sri Lanka Tall, Sri Lanka Green Dwarf, Bodiri Tall, San Ramon, King coconut and Sri Lanka Brown dwarf. The information gathered in the morphological characterization was used to classify the new coconut phenotypes into the existing taxonomic classification of coconut in Sri Lanka. Accordingly, the new phenotypes, Ran pol, Juwan, Thatin, Dothalu and Rath gon thembili were classified as different forms within the variety *Typica*, Murusi under variety *Nana* and Bothal thembili within the variety *Aurantiaca*. A higher level of morphological diversity was observed and the multivariate data analysis methods revealed the groupings of the new phenotypes along with their comparative coconut forms.

Molecular characterization was performed on 40 genotypes of coconut representing the new coconut phenotypes and their comparative coconut forms, at 13 microsatellite marker loci. All the markers were observed to be polymorphic and thus informative and a total of 88 different alleles were scored at a mean allele frequency of 6.7692 and a mean gene diversity of 0.6784. The genetic distances and the phenetic tree derived using genotypic data revealed the genetic relationships, genetic relatedness and the genetic diversity among the new coconut phenotypes and between their comparative coconut forms. The new coconut forms identified, characterized and conserved in this research will enhance the breeders' collection of coconut.