

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317178669>

Impact of routine laboratory culture media on in-vitro biofilm formation of *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Enterococcus faecalis*

Conference Paper · May 2017

CITATIONS

0

READS

61

8 authors, including:



Gayan Kanchana Wijesinghe

University of Campinas

11 PUBLICATIONS 10 CITATIONS

[SEE PROFILE](#)



Ayomi Dilhari

University of Sri Jayewardenepura

14 PUBLICATIONS 14 CITATIONS

[SEE PROFILE](#)



Gd Buddhika Gayani

University of Sri Jayewardenepura

7 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Chinthika Gunasekara

University of Sri Jayewardenepura

66 PUBLICATIONS 125 CITATIONS

[SEE PROFILE](#)


Some of the authors of this publication are also working on these related projects:



Effect of Citrate encapsulated LDH, Curcumin intercalated LDH and other modified nano-particles on in-vitro *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Enterococcus faecalis* biofilms. [View project](#)



Desalination of Sea and Hard water [View project](#)



The SCIENTIFIC SESSIONS 2017

“Building Bridges for Better Health”

Faculty of Medical Sciences, University of Sri Jayewardenepura
in collaboration with
Colombo South Teaching Hospital
Sri Jayewardenepura General Hospital
Base Hospital, Homagama

Book of Proceedings

24th - 25th May 2017



OP 9

Impact of routine laboratory culture media on *in-vitro* biofilm formation of *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Enterococcus faecalis*

Wijesinghe GK¹, Dilhari KAA¹, Buddhika G², Gunasekera TDCP¹, Fernando SSN¹, Kottegoda N², Samaranyake LP³, Weerasekera MM¹

¹Department of Microbiology, Faculty of Medical Sciences, University of Sri Jayewardenepura, ²Department of Chemistry, Faculty of Applied Sciences, University of Sri Jayewardenepura ³Health Science Center, Kuwait University, Kuwait City, Kuwait

Objectives: This study was aimed to determine the efficacy of four routine laboratory culture media on biofilm formation of *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Enterococcus faecalis*.

Methods: A sterile flat bottom 96 well plate was inoculated using 0.5 McFarland equivalent standard cell suspension of *P. aeruginosa*, *S. aureus* and *E. faecalis* and the growth rate of planktonic cells was quantified by measuring the optical density (OD₄₉₂) at two hour intervals. Influence of culture medium on adhesion of bacteria as an initial step of biofilm formation in the presence of four culture media

(Nutrient broth (NB), Brain Heart Infusion (BHI) broth, Luria-Bertani (LB) broth and RPMI 1640) was quantified using MTT (3-[4, 5- dimethylthiazole-2-yl]-2, 5- diphenyltetrazolium bromide) assay after 90 minutes adhesion. Biofilms of *P. aeruginosa*, *S. aureus*, *E. faecalis* and their 1:1 mixed biofilms were developed and the growth was quantified using MTT metabolic activity at 24 hour time intervals. Scanning electron microscopy (SEM) was performed to assess the ultrastructure.

Results: On comparing the relative growth of the bacteria in different culture media, the maximum growth of all three planktonic cultures was achieved using BHI broth. All mono species and mixed species cultures exhibited their maximum adhesion in the presence of RPMI 1640. All biofilm exhibited the maximum growth in BHI broth. SEM imaging had shown the enhanced growth of ultrastructure of the biofilm with the presence of BHI broth.

Conclusions: The maximum planktonic and biofilm growth was achieved with BHI broth. However, bacterial adhesion was enhanced in the presence of RPMI 1640.