



Biochemical changes during ripening of banana: A review

*SDT Maduwanthi, RAUJ Marapana

Department of Food Science and Technology, Faculty of Applied Sciences, University of Sri Jayewardenepura, Sri Lanka

Abstract

In fruit ripening fruits undergo different biochemical and physical changes and those changes are characteristic to different fruits. Studying the nature of those changes may be beneficial for postharvest handling of particular fruits. The aim of this review is to study biochemical changes that occur during ripening process in banana. During ripening process softening of fruit flesh occurs mainly due to degradation of starch and insoluble polysaccharides. As well, sugar content is increased from 2% to 20% during ripening. Colour changes of banana peel occur due to degradation of chlorophylls. Malic acid content in flesh is said to increase during ripening process. Astringency, which occurs due to tannin, is decreasing during ripening. Volatile profile undergoes different changes in different changes of the ripening process.

Keywords: banana, ripening, biochemical changes

1. Introduction

Banana belongs to the genus *Musa* of the family *Musaceae*. The genus *Musa* contains four sections, *Eumusa*, *Rhodochlamys*, *Australimusa*, and *Callimusa*. *Rhodochlamys* and *Callimusa* are of ornamental interest only. *Australimusa* species are utilized across a large area of the Pacific as a cooked vegetable. However, *Eumusa* is the largest and most widespread geographically and contains all the major edible species of banana. Most edible bananas are derived from two members of the sect of *Eumusa*: *Musa acuminata* and *Musa balbisiana* [1]. Most of the edible bananas are either derived solely from the *Musa acuminata* or are hybrid between two wild diploid species, *Musa acuminata* Colla and *Musa balbisiana* Colla; which contributed to A and B genomes, respectively [2].

There are 300-500 cultivars which are naturally occurring banana hybrids which can be grouped according to their genome constitutions, as AA, AAA, AB, AAB, ABB and ABBB. Tetraploid AAAA cultivars have only been produced through breeding program. Nevertheless triploid and tetraploid B cultivars have not been identified yet [3]. Three most common species of banana which are mostly grown in the world are *Musa cavendishi*, *Musa paradisiaca* and *Musa sapientum* [4].

It is one of the top most fruit in the world in production and as well as in consumption. Bananas are climacteric fruits which are artificially ripened regularly. Ripening process of banana can be divided into three distinct phases namely the pre-climacteric or 'green life' stage, the climacteric and ripening stage and finally eat-ripe and senescence stage [5]. During ripening process banana undergoes different physiological, biochemical, and organoleptic changes that lead to a soft and edible ripe fruit [6]. This review paper investigates the changes that occur during banana ripening.

2. Bio chemical changes during ripening

Fruit ripening is a genetically programmed, highly coordinated process of organ transformation from unripe to ripe stage, to yield an attractive edible fruit with an optimum blend of colour, taste, aroma and texture. A set of biochemical and physical changes occur in banana during ripening which makes it an edible fruit. These changes involve several biochemical pathways like degradation of starch to sugar, change in the peel and pulp colour, cell wall changes, change in the concentration of volatiles and acids [6].

2.1 Tissue softening

Softening is a very important aspect of the ripening syndrome. Loss of turgor, degradation of starch and enzyme catalyzed changes to wall structure and composition are the mechanisms which lead to fruit softening. According to Finney [7] textural change in banana fruit during ripening is predominantly due to the changes in chemical structure of starch grains. Many researchers have shown that starch content in the pulp of banana decreases drastically during the short period of ripening and then starch is no longer detected [8, 9, 10]. However, Kojima [11] suggests that banana pulp softening process is due to the associated process whereby the contents of pectic and hemicellulosic polysaccharides and starch decrease during ripening. According to ALI [12] 50% firmness loss was occurred in 3 days during ripening in Mas banana (*Musa acuminata*, AA group).

2.2 Carbohydrates

During ripening process starch is converted into simple sugars through enzymatic browning process [13]. Starch forms about 20 to 25% of the fresh weight of the pulp of unripe bananas. Sugars are present in the green fruit only about 1 to 2% in the fresh pulp which rises up to 15 to 20% at ripeness [14]. The soluble sugars detected in ripened banana are mainly

sucrose, glucose and fructose [15]. According to Adao and Gloria [15] the mean level of starch content in 'Prata' banana was reduced from 15.7 g/100 g to 3.40 g/100 g during ripening. As well total soluble sugar content was increased from 1.26g/100g to 14.3g/100g. Adewale [16] reported that unripe banana (*Musa sapientum*) had highest amylase activity (3900 ± 310 Units/mg protein) and decreased rapidly to a very low value (100 ± 15 Units/mg protein) when it was fully ripened.

2.3 Pigments

The peel colour changes from green to yellow during ripening of banana fruit. The most important compounds responsible for the change in peel color are chlorophylls and carotenoids [17]. Chlorophyll content decreases become absent in ripe fruit [18, 19, 20]. The level of total carotenoids decreased to half the level at the colour break and subsequently again reached a level similar to that in green fruit [18]. According [21] the major pattern of pulp carotenoids is α -carotene (31%), β -carotene (28%) and lutein (33%) of total carotenoids. Banana peel contained 3-4 $\mu\text{g/g}$ carotenoids content as lutein equivalent and ingredients were lutein, β -carotene, α -carotene, violaxanthin, auroxanthin, neoxanthin, isolutein, β -cryptoxanthin and α -cryptoxanthin.

2.4 Pectin

The inter-lamella layer in higher plants composed of polysaccharides mainly pectin [22]. The nature of the pectic component in cell wall is associated with fruit softening [5]. Pectin is a linear chain of α -(1 \rightarrow 4)-linked D-galacturonic acid which are methyl esterified. The great strength of green fruit is due to the protopectin or water insoluble pectic which is partially esterified polygalacturonic acid [23]. Increased solubility of pectic polysaccharides is one of the identified changes happens during fruit ripening. Pectic enzymes are related with the softening of fruits along with the increase in soluble pectins. The mainly enzymes involved in pectin degradation are Polygalacturonase (PG) and pectin methylesterase (PME). According to Tapre and Jain [24] pectin content in banana (*Musa sp.* var 'Robusta') pulp increased from 0.37-0.66% significantly which was measured as calcium pectate. According to Robinson and Saucó [5] reported ripe pulp in banana contains 0.5-0.7% pectin content. Whereas several studies have been done to characterize pectin degrading enzymes and their activity. Smith [25] and Markovic [26] reported the presence PG in banana tissue. Nagel and Patterson [27] identified that PME activity appears as the fruit ripens and continues to increase exponentially. However Patil and Magar [28] reported that PME activity is highest at colour stage 4 (greenish yellow) and fell down sharply in the advanced stages of ripening.

2.5 Organic acids

Vonloesecke [29] reported the presence of malic, citric, oxalic, and tartaric acids in the banana fruit, with malic acid being the principal acid. Malic and Citric acids are responsible for tartness in the unripe banana while oxalic acid is contributed to astringent taste of the fruit [19]. According to von loesecke [29] titratable acidity of the pulp increases to a peak during ripening in the case of some varieties, and then declines.

Further it has reported that malic acid concentration has been reported to vary between 0.8 and 7.5 meq/100g and it is increasing three to sevenfold during ripening [29]. The controversial results with Vonloesecke [29] were reported in Soltani [30] where titratable acidity decreased gradually until the fruit reaches to full-ripe (stage-6) then increased at stage-7. In Malakar [31] and Kulkarni [32], titratable acidity was decreased during the ripening period.

According to Wyman and Palmer [33] malic acid content was 1.36 meq/100g at the pre climacteric stage and it increased up to 5.37 at the climacteric stage and 6.2 meq/100g at the post climacteric stage while oxalic acid content was decreased from 2.33 to 1.37 meq/100g at pre climacteric stage to post climacteric stage. However total organic acidity was increased during ripening.

2.6 Astringency

Most fruits and vegetables are astringent in unripe stage and during ripening process astringency is reduced. Similarly banana which tastes astringent in unripe stage becomes palatable in the eating ripe stage. Astringency is claimed to be related to tannins which causes drying, roughing and puckering of the mouth epithelia giving astringency in oral sensation [34]. Unripe fruit contained considerable level of tannin which reduces as ripening proceeds [29, 35]. However some earlier researches reported that tannin remained unchanged during ripening. The confusion of reduction of astringency and constant value of tannin content was well explained by Barnell and Barnell [34]. According to Von loesecke [29] there are many types of tannin and during ripening soluble tannin becomes insoluble. Barnell and Barnell [34] explained that there are two types of tannin containing elements, latex vessels in both the pulp and peel and small scattered cells in the peel. Further Barnell and Barnell [34] explained that nature of tannin in latex vessels undergo changes while tannin content of the small scattered cells of the skin appears to undergo little or no change.

2.7 Volatile constituents

Very few researches have been published on aroma compounds and their changes during ripening. The unique aroma of bananas arises from set of volatile constituents including esters, alcohols, ketones, aldehydes and phenol esters [19]. According to Pino and Febles [36]. The composition of banana (*Musa spp.*, AAA group) fruit volatiles included 75 esters, 18 ketones, 14 phenols and derivatives, 7 aldehydes, 13 alcohols, 7 acids, and 12 miscellaneous compounds. As well isoamyl and isobutyl esters together with 2-pentanone are the major compounds found in banana volatile profile [37]. Mccarthy [38] reported that characteristic banana flavour is due to the amyl esters of acetic, propionic and butyric acids. Jordan [37] studied aromatic profile of fresh banana fruit paste and identified 26 volatile compounds including *E*-2-hexenal and hexanal as major aldehydes and 3-hydroxy-2-butanone, 3-methyl-1-butanol, 2-pentanol, isoamyl acetate, isoamyl isobutyrate, and eugenol were also presented in high concentrations.

3. Conclusion

Ripening is a process consists of a set of biochemical and

physical changes which gives an edible fruit. Banana is one of mostly consumed fruit in the world and it has been studied for its biochemical and physical changes for years. Softening of the texture, yellowing of peel, reduction of astringency and increase of sweetness are major organoleptic changes which can be noted in banana ripening. These changes occur as a result of series of biochemical changes in peel and flesh of banana fruit.

4. References

- Salunkhe DK, Kadam SS. Handbook of Vegetable Science and Technology: Production, Compostion, Storage, and Processing. CRC press, 1998, 19.
- Hailu M, Workneh TS, Belew D. Review on postharvest technology of banana fruit. African Journal of Biotechnology, 2013, 12(7).
- Hirimburagama K, Gamage N. In vitro multiplication of local cultivars of banana (*MUSA* spp.) through shoot-tip culture. Journal of the National Science Foundation of Sri Lanka, 2015, 24(1).
- Mohapatra D, Mishra S, Sutar N. Banana and its by-product utilisation: an overview. Journal of Scientific and Industrial Research. 2010; 69:323-329.
- Robinson JC, Saúco VG. Bananas and plantains, 2010.
- Gowen S. Bananas and plantains. Springer Netherland. 1995.
- Finney EE, BEN-GERA IT, MASSIE DR. An objective evaluation of changes in firmness of ripening bananas using a sonic technique. Journal of food science. 1967; 32(6):642-6.
- Barnell HR. Studies in Tropical Fruits: XV. Hemicellulose Metabolism of the Banana Fruit during Storage and Ripening. Annals of Botany. 1943; 7(28):297-323.
- Charles RJ, Tung MA. Physical, rheological and chemical properties of bananas during ripening. Journal of food science. 1973; 38(3):456-9.
- Agravante JU, Matsui T, Kitagawa H. Starch breakdown in ethylene-and ethanol-treated bananas: Changes in phosphorylase and invertase activities during ripening. Nippon Shokuhin Kogyo Gakkaishi. 1990; 37(11):911-5.
- Kojima K. Softening of banana fruit- Relationship between firmness and chemical composition. Japan Agricultural Research Quarterly. 1996; 30:269-274.
- Ali ZM, Chin LH, Lazan H. A comparative study on wall degrading enzymes, pectin modifications and softening during ripening of selected tropical fruits. Plant Science. 2004; 167(2):317-27.
- Yang SF, Hoffman NE. Ethylene biosynthesis and its regulation in higher plants. Annual review of plant physiology. 1984; 35(1):155-89.
- Simmonds NW. Bananas. Webster printing service Ltd Bristol. Imperial College of Tropical Agriculture. Trop. Agric, 1959, 453-61.
- Adão RC, Glória MB. Bioactive amines and carbohydrate changes during ripening of Prata banana (*Musa acuminata* × *M. balbisiana*). Food Chemistry. 2005; 90(4):705-11.
- Adewale IO, Adefila A, Adewale TB. Changes in Amylase Activity, Soluble Sugars and Proteins of Unripe Banana and Plantain during Ripening. Annual Review & Research in Biology. 2013; 34:678-685.
- Subagio A, Morita N, Sawada S. Carotenoids and their fatty-acid esters in banana peel. Journal of nutritional science and vitaminology. 1996; 42(6):553-66.
- Gross J, Flugel M. Pigment changes in peel of the ripening banana (*Musa cavendishi*). Gartenbauwissenschaft. 1982; 47(2):62-4.
- Seymour GB, Thompson AK, John P. Inhibition of degreening in the peel of bananas ripened at tropical temperatures. Annals of applied biology. 1987; 110(1):145-51.
- Marriott J, Palmer JK. Bananas-physiology and biochemistry of storage and ripening for optimum quality. Critical Reviews in Food Science & Nutrition. 1980; 13(1):41-88.
- Gross J, Carmon M, Lifshitz A, Costes C. Carotenoids of banana pulp, peel and leaves. Food Science and Technology. 1976; 9:211-4.
- Verma C, Singh RK, Mishra S. Biochemical Characterization of Pectin Methylsterase from *Musa acuminata* Referring to Delayed Ripening. IOSR Journal of Pharmacy and Biological Sciences. 2014; 9:42-47.
- Joslyn MA. The chemistry of protopectin: a critical review of historical data and recent developments. Advances in food research. 1963; 11:1-07.
- Tapre AR, Jain RK. Study of advanced maturity stages of banana. International Journal of Advanced Engineering Research and Studies. 2012; 1(3):272-274.
- Smith NJ, Seymour GB, Jeger MJ, Tucker GA. Cell wall changes in bananas and plantains. In Symposium on Tropical Fruit in International Trade 269, 1989, 283-290.
- Markovič O, Heinrichova K, Lenkey B. Pectolytic enzymes from banana. Collection of Czechoslovak Chemical Communications. 1975; 40(3):769-74.
- Nagel CW, Patterson ME. Pectic enzymes and development of the pear (*Pyrus communis*). Journal of Food Science. 1967; 32(3):294-7.
- Patil DL, Magar NG. Correlation between banana fruits ripening, its texture and pectin-methyl esterase activity. In: Proceedings of the Indian Academy of Sciences-Section B. 1975; 81(3):127-130.
- Von Loesecke HW. Bananas: chemistry, physiology, technology. Interscience Publishers, Inc, 1950.
- Soltani M, Alimardani R, Omid M. Comparison of some chromatic, mechanical and chemical properties of banana fruit at different stages of ripeness. Modern Applied Science. 2010; 4(7):34.
- Malakar S, Roy S, Kumar N. Evaluation of Changes in Physico-Chemical Parameters of Banana (*Musa Spp.*) during Ripening in Storage. Journal of Agricultural, Engineering and Food Technology. 2015; 2:100-103.
- Kulkarni SG, Kudachikar VB, Prakash MK. Studies on physico-chemical changes during artificial ripening of banana (*Musa sp*) variety 'Robusta'. Journal of Food Science Technology 2011; 48(6):730-734.
- Wyman H, Palmer JK. Organic acids in the ripening banana fruit. Plant physiology. 1964; 39(4):630
- Barnell HR, Barnell E. Studies in tropical fruits: XVI. The distribution of tannins within the banana and the

- changes in their condition and amount during ripening. *Annals of Botany*. 1945; 9(33):77-99.
35. Slocum AH. Banana Chilling Studies. II. Effect of Low Temperature upon Cell Structure. *Bull*, 1933, (48).
 36. Pino JA, Febles Y. Odour-active compounds in banana fruit cv. Giant Cavendish. *Food chemistry*. 2013; 141(2):795-801.
 37. Jordán MJ, Tandon K, Shaw PE, Goodner KL. Aromatic Profile of Aqueous Banana Essence and Banana Fruit by Gas Chromatography- Mass Spectrometry (GC-MS) and Gas Chromatography- Olfactometry (GC-O). *Journal of agricultural and food chemistry*. 2001; 49(10):4813-7.
 38. McCarthy AI, Palmer JK, Shaw CP, Anderson EE. Correlation of Gas Chromatographic Data with Flavor Profiles of Fresh Banana Fruit. *Journal of Food Science*. 1963; 28(4):379-384.